

Average Cosmological Invariant Parameters of Cosmic Gamma Ray Bursts

Igor G. Mitrofanov¹, Dmitrij S. Anfimov¹, Maxim L. Litvak¹

and

Michael S. Briggs², William S. Paciesas², Geoffrey N. Pendleton², Robert D. Preece²

and

Charles A. Meegan³

¹Space Research Institute, Profsojuznaya str. 84/32, 117810 Moscow, Russia

²Department of Physics, University of Alabama in Huntsville, Huntsville, AL 35899

³NASA/Marshall Space Flight Center, Huntsville, AL 35812

Received _____; accepted _____

ABSTRACT

Average cosmological invariant parameters (ACIPs) are calculated for six groups of BATSE cosmic gamma-ray bursts selected by their peak fluxes on the 1.024 s time scale. The ACIPs represent the average temporal and spectral properties of these events equally in the observer frame of reference and in the co-moving frames of outbursting emitters. The parameters are determined separately for rise fronts and for back slopes of bursts, defined as the time profiles before and after the main peaks, respectively. The ACIPs for the rise fronts are found to be different for different intensity groups, while the ACIPs for the back slopes show no significant dependence on intensity. We conclude that emitters of bursts manifest standard average properties only during the back slopes of bursts.

1. Introduction

The isotropy of gamma-ray bursts on the sky combined with the significant deviation of the $\log N/\log P$ curve from a $-3/2$ power law (Meegan et al. 1992) gave the first clear evidence that sources of gamma-ray bursts are at cosmological distances, where the deficit of observed number of dim events is a direct consequence of the non-Euclidean-nature of the expanding Universe (Paczynski 1986). Strong evidence in favor of the cosmological paradigm has recently been provided by the detection of red-shifted spectral lines from the optical counterparts of GRB 970508 (Metzger et al. 1997) and GRB 971214 (Kulkarni et al. 1998).

As soon as the cosmological paradigm became popular for gamma-ray bursts, two effects were suggested to test it using the observational data of gamma-ray bursts. The first one is the effect of cosmological red shift, which predicts that the energy of spectral features of dimmer bursts should be red-shifted in respect to similar features of bright events. The second one is the complementary effect of cosmological time dilation, which predicts that light curves of dimmer bursts should be time-stretched with respect to those of brighter events.

Since both effects are associated with the geometry of the expending Universe, we will refer to them as *geometrical* effects. The transformation factors are known to be

$$Y(z_{\text{br}}, z_{\text{dim}}) = (1 + z_{\text{dim}})/(1 + z_{\text{br}}), \quad (1)$$

where z_{br} and z_{dim} are the red shifts of emitters of bright and dim bursts, respectively.

If the brightest bursts are associated with the nearest objects and the dimmest bursts are related to the furthest emitters, their intensities could be used to determine distances to the corresponding emitters. In this case the cosmological effects should lead to a hardness/intensity correlation (due to the red shift), and a stretching/intensity

anti-correlation (due to the time dilation).

However, gamma-ray bursts are known to be very different from each other. In searching for the generic effects of hardness/intensity correlations and stretching/intensity anti-correlation between different intensity groups, one has to identify generic signatures among the divergence of properties of individual events to represent the properties of *typical* burst emission.

Using such signatures, the effects of hardness/intensity correlation and stretching/intensity anti-correlation have been studied by comparing groups of BATSE bursts with different brightnesses. The average peak energy E_p of νF_ν spectra was found to show a hardness/intensity correlation consistent with cosmological red shift (Mallozzi et al. 1995), and there is evidence that average emissivity curves of dimmer bursts are stretched relative to those of brighter bursts, although the results of different analyses are not entirely consistent (Norris et al. 1994, Norris et al. 1998, Mitrofanov et al. 1996, Mitrofanov et al. 1998). The possibility that the stretching of dimmer bursts might result from a selection effect has been considered, and it has been shown that it is not a selection effect (Wijers & Paczynski 1994). On the other hand, it is also not clear that the results of red shift and time dilation studies agree with each other (Mitrofanov 1996).

On the other hand, the logically simple concept that burst intensities are “standard candles” may not be correct. It has been shown (Brainerd 1997) that the broad spread of observed intensities of bursts could result from a broad spread of intrinsic luminosities of emitters. Also, three bursts with measured red-shifts of afterglowing optical emission have gamma-ray luminosity differing over a range of ~ 20 times. Despite of small sample size, these three events provide quite good evidence that GRBs are not standard candles. The correlations of burst hardness and duration with intensity could be related in this case to the intrinsic properties of sources rather than to geometrical effects of cosmological space

(citeMit2). Furthermore, one should take into account that the astronomical population of emitters could be quite different at close and distant cosmological distances. In this case the difference between brightness groups could reflect the intrinsic evolution of sources rather than cosmological effects.

Therefore, to test the cosmological paradigm it is necessary to separate in the data the *geometrical effects* due to cosmological expansion of the Universe from the *physical effects* due to intrinsic luminosity-based correlation of emitters and the *astronomical effects* due to evolution of sources over the different red shifts.

2. Double-peak energy, as a burst spectral parameter

To compare the average temporal and spectral signatures of gamma-ray bursts with different brightness, 6 intensity groups, each consisting of ~ 100 bursts, were selected by their peak fluxes F_{max} on the 1024 ms time scale in the 3B BATSE Catalog (Meegan et al. 1996), as shown in Table 1.

The time profile of each burst has a well-defined moment t_{max} when the observed flux reaches a maximum F_{max} . One might postulate that peaks are associated with some particular physical transition, when the average rising trend of intensity before the peak is converted into the decaying tail after that. The procedure of peak alignment has the physical sense to combine all bursts together at the same stage of the emission process. On the other hand, the νF_ν energy spectrum of burst emission usually has a well-defined peak energy E_p , but E_p typically varies with time during a burst. One could, however, introduce a single-valued spectral parameter for each burst, which is the peak energy $E_{p,\text{max}}$ of the spectrum at the time of peak photon flux F_{max} . We will refer to $E_{p,\text{max}}$ as the *double-peak energy*, which corresponds to photons with the largest spectral density of emission.

We evaluated the double-peak energy $E_{p,\max}$ for each BATSE burst using the CONT data, which has a time resolution of 2048 ms (Anfimov et al. 1998). The distributions of these energies for the reference group and for the dimmest intensity group are shown in Figure 1. The observed distributions for these groups might be compared with K-S test provided that the reference group is shifted leftward to match the curves for the dimmer groups. The lowest K-S probability is equal to 0.3 for the comparison between reference group and dimmest intensity group when they are shifted to the same mean. The present data allows us to measure with statistical significance these shifts, but does not reveal any difference in the shapes. We might assume that there is an *universal* log-normal law for all these distributions, which can be parameterized using a set of red-shifting factors $Y^{(i)}$, defined by shifting the log-normal distribution for the brightest reference group ($i = 1$) to provide the best fit for the corresponding distributions of the dimmer intensity groups ($i = 2-6$).

Table 1 shows the best-fitting shift factors and the log-normal average values $E^{(i)} = \langle E_{p,\max}^{(i)} \rangle$ for our selected intensity groups. The differences between them show the effect of hardness/intensity correlation of gamma-ray bursts. Basically, the ratios between the average double-peak energies for different brightness groups $E^{(1)}/E^{(i)}$ are very close to the best fitting shifting factors $Y^{(i)}$. The difference of the average peak fluxes between the 100 brightest and 100 dimmest bursts is a factor of ~ 43 , while the corresponding log-normal average values of the double peak energy differ by a factor of ~ 3 (Table 1).

3. Equivalent time width, as a burst temporal parameter

A robust temporal parameter for bursts is very difficult to define. Bursts have a variety of light curves, and for many of them the light curves are very complex. The evaluation of a *duration-type* parameter depends on the sensitivity of the instrument, its energy range,

and its time resolution. The best known parameters t_{50} and t_{90} attribute definite durations to any individual burst, but there are several statistical biases that could affect the results when statistical studies are performed using them.

Below we suggest another duration-type parameter that could be associated with a group of bursts. It was defined using the Average Emissivity Curves (Mitrofanov et al. 1996). The ACE is known to be a quite robust signature for any large group of bursts. This ACE profile represents the slow component of burst variability, which can be interpreted as the *general envelope* of individual light curves. By its nature, the ACE averages over the faster variability, leaving only the signature of *slow clocks*, so in computing the ACE of BATSE gamma-ray bursts we can use the DISCLA and CONT data, which have 1024 ms and 2048 ms time resolution, respectively. The ACE intermixes individual events with all their particular time profiles, and represents them by one profile, which is a single asymmetric peak with steeper rise front and flatter back slope. Therefore, for each intensity group i the ACE profile can be used to estimate the average duration of rise fronts and back slopes for averaged events.

Although the BATSE data set is large, the variety of burst time profiles is such that we must consider its effect on the variance among finite samples. We have found by direct comparison of ACE profiles for different samples of BATSE bursts that they were much more distinct than would be expected from the errors of sample variance for individual samples. This means that a random sample of bursts for individual groups does not ensure the well-weighted contribution of events with all kinds of profiles. To study the random choice statistics of bursts, a special Monte Carlo simulation was performed using the total set of 603 bursts in the 3B Catalog with durations $t_{90} > 2$ s (Mitrofanov et al. 1998, Litvak et al. 1998). Indeed, the distributions of stretching coefficients Y due to statistics of random choice were found to be much broader than expected from the sample variance predicted

by normal statistics. From the simulations, the 1σ deviation around a non-stretched value $Y \sim 1$ for a sample of N bursts was found to be

$$\frac{\delta Y}{Y} = 0.13 \cdot \sqrt{\frac{100}{N}}. \quad (2)$$

This value can be used as an estimate of the error in stretching factors between ACEs for any two groups of N bursts. The present size of our selected intensity groups (Table 1) allows us to resolve stretching at $\sim 3\sigma$ significance between them provided the effect is larger than ~ 1.41 (Mitrofanov et al. 1998, Litvak et al. 1998).

There is a simple analytic form that provides a very good fit to the ACE profile $\Phi(t)$ for time intervals of 20–50 s around the maximum:

$$\Phi(t) = \left(\frac{t_0}{t_0 + |t - t_{\max}|} \right)^a, \quad (3)$$

where the exponent a has different values a_{RF} and a_{BS} for the rise front and back slope, respectively. This law allows us to take into account the energy dependence of ACEs (Mitrofanov et al. 1997b) by interpolating in energy between the parameters a_{RF} , a_{BS} and t_0 measured for ACEs in three BATSE discriminator channels (25–50 keV, 50–100 keV and 100–300 keV). It also allows us to use a time-efficient procedure to estimate the relative time-stretching factor between ACE profiles for any two samples with different intensity (Mitrofanov et al. 1998).

The procedure to build the ACE includes the selection of the highest peak of each burst C_{max} *in count space* and the normalization of time profiles by the C_{max} value. Therefore, the ACE is sensitive to a bias resulting from domination of positive fluctuations in the selected peaks. Due to this bias the ACE profile is systematically lower in both wings; i.e., the measured value is narrower than the true value. The bias is stronger for dimmer bursts, where the influence of positive fluctuations is larger. To take it into account, our reference group 1 (the brightest one) was transformed into an artificial reference group by Monte

Carlo noisification, having the same event profiles but with dimmer peak fluxes. When the reference group 1 was dimmed down to the level of the dimmest group 6, the noise-produced ACE was found to be different than the original ACE by a factor of ~ 0.8 (that is: narrower – see Mitrofanov et al. 1998).

To estimate the average equivalent time width for the brightest reference group 1, the observed ACE profiles in three energy ranges $j = 1$ (25-50 keV), $j = 2$ (50-100 keV) and $j = 3$ (100-300 keV) were used as they are. The bias due to positive fluctuations is assumed to have no influence on this group. Using the $\text{ACE}^{(1,j)}$ profiles, we calculated average equivalent widths in each of three energy channels j as

$$\tau_{\text{RF,BS}}^{(1,j)} = \int dt \Phi_{\text{FR,BS}}^{(1,j)}(t) \quad (4)$$

for the rise front (RF) and back slope (BS), respectively. The values of $\tau^{(1,j)}$ were then interpolated over the broad energy range 25–300 keV, and the equivalent width was determined at the double-peak energy $E^{(1)} = 293$ keV. The values of $\tau^{(1,j)}(E^{(1)})$ for rise front and back slope are the temporal parameters $\tau_{\text{RF,BS}}^{(1)}$ for the reference group 1 (Table 1). Physically, τ represents the average duration of emission either over the rise or over the decay at the spectral range around the double-peak energy $E_{\text{p,max}}$.

$\text{ACE}^{(i,j)}$ profiles for dimmer groups ($i = 2-6$) can be used similarly to estimate temporal parameters, provided that they are corrected for the noise-produced narrowing of ACEs. For a given intensity group i an artificial reference group i' was created from the events of the reference group 1 by Monte Carlo noisification, in which the reference bursts are reduced in intensity to the fluxes of group i , and noise added corresponding to the noise level of group i . The artificial reference group i' therefore represents the original group 1, but takes into account the noise-produced effects. Therefore, to evaluate the noise-corrected average stretching between the testing group i and the reference group 1 for energy channel j , we measured the stretching factors $Y^{(i,j)}$ between the $\text{ACE}^{(i,j)}$ of the actual dim group

i and the $\text{ACE}^{(i',j)}$ of the artificial reference group i' . We calculated $Y^{(i,j)}$ separately for rise fronts (RF) and back slopes (BS) for intensity groups $i = 2\text{--}6$ in three energy ranges $j = 1\text{--}3$. Using these factors, the parameters of equivalent width of $\text{ACE}^{(i,j)}$ could be defined for the energy channels j , as

$$\tau_{\text{RF,BS}}^{(i,j)} = \tau_{\text{RF,BS}}^{(1,j)} \cdot Y_{\text{RF,BS}}^{(i,j)}. \quad (5)$$

These values are corrected for the noise-produced narrowing of ACEs because they are determined from the stretching factors between the testing dim groups and the corresponding artificially noisified reference groups. The single-value temporal parameters $\tau^{(i)}(E^{(i)}) = \tau^{(i)}$ were interpolated between the values $\tau_{\text{RF,BS}}^{(i,j)}$ for three discriminator channels $j = 1\text{--}3$ at the double-peak energies $E_{\text{p,max}}^{(i)}$. The parameters $\tau^{(i)}$ are presented in Table 1 for rise fronts (RF) and back slopes (BS) of bursts. Their errors are estimated from the choice statistics for stretching factors $Y_{i,j}$. The rise front equivalent widths $\tau_{\text{RF}}^{(i)}$ do not show a correlation with burst intensity. On the other hand, the back slope equivalent widths $\tau_{\text{BS}}^{(i)}$ are significantly increasing with decreasing intensity of bursts.

4. Average Cosmological Invariant Parameters

The average double-peak energy $E_{\text{p,max}}^{(i)}$ and the equivalent time width $\tau^{(i)}$ at the double-peak energy are very useful parameters for testing the cosmological paradigm. Indeed, the energy-dimension parameters $E_{\text{p,max}}^{(i)}$ represent *spectral signatures* that have the same physical sense for all bright, medium and dim groups of bursts. The time-dimension parameters $\tau^{(i)}$ represent *temporal signatures* that are also well-defined for all groups of bursts.

We have found that these parameters vary significantly among the different intensity groups. However, the differences between them could be caused either by the purely

geometrical transformations of red shift and time dilation in the expanding Universe, or by a *physical* variation among the outbursting sources in the co-moving frames. These two parameters cannot by themselves be used to perform a model-independent test of the cosmological paradigm of gamma-ray bursts. One must either postulate some intrinsic properties of emitters and then resolve the cosmological transformations of observed gamma-ray bursts, or postulate the geometrical effects of time dilation and energy red shift and then deconvolve properties of observed bursts into the intrinsic properties of emitters.

As an alternative, we wish to find a special observational parameter for any selected sample of gamma-ray bursts that does not depend on the geometrical effects of the Universe extension, and which we call an *Average Cosmological Invariant Parameter* (ACIP)(Mitrofanov et al. 1998a). Let us assume that some brightness group i corresponds to emitters with red shifts around some average value $z^{(i)}$ and the corresponding equivalent width and double-peak energy equal $\tau_*^{(i)}$ and $E_*^{(i)}$ in the co-moving frame, respectively. Then, since a time-dimensional average parameter is increased by a factor $(1 + z^{(i)})$, giving $\tau^{(i)} = \tau_*^{(i)}(1 + z^{(i)})$ in the observer's frame, and an energy-dimensional average parameter is reduced by a factor $(1 + z^{(i)})^{-1}$, giving $E^{(i)} = E_*^{(i)}(1 + z^{(i)})^{-1}$ in the observer's frame, the product $\Pi^{(i)}$ of time-dimensional and energy-dimensional average parameters for the group is an invariant because the red shift factors cancel each other:

$$\Pi^{(i)} = \tau^{(i)} \cdot E^{(i)} = \tau_*^{(i)} \cdot E_*^{(i)}. \quad (6)$$

Therefore, any difference between values of Π for two different samples of bursts has to be attributed to a real physical difference between their emitters.

5. Results from a comparison of ACIPs for different intensity groups

The average durations of rise fronts and back slopes are known to have different behaviors for bursts with the different intensity. We calculated Π as defined in equation (6) for the same 6 brightness groups as in Table 1, separately for time signatures τ_{RF} and τ_{BS} . The results are presented in Table 2 and Figures 2 & 3.

One can consider emitters of gamma-ray bursts as *standard candles* with respect to the property described by the ACIP if the values are independent of brightness. This model can be rejected for the rise fronts of bursts: the assumption of a constant Π_{RF} in our analysis has a negligibly small probability (< 0.001). Indeed, Π_{RF} decreases with decreasing average fluxes $\langle F_{\text{max}} \rangle$, as

$$\Pi_{\text{RF}} = 1196 \cdot \left[\frac{\langle F_{\text{max}} \rangle}{14.2} \right]^{0.22 \pm 0.03}, \quad (7)$$

where $\langle F_{\text{max}} \rangle$ is in units of $\gamma/\text{cm}^2\text{-s}$ and Π_{RF} is in units of keV-s. Therefore Π_{RF} cannot be used as a *standard candle* (Figure 2). Of course, there may be a concern that the dimmest group suffers from incompleteness, since it is closest to the trigger threshold where slow-rising events can be missed (Kommers et al. (1997)). If we exclude the dimmest group from the fit in Figure 2, the fitted power-law index in eq. 7 does not change dramatically: 0.18 ± 0.03 . In this case, the probability that the data are consistent with a constant is still very small (0.003).

On the other hand, the values of Π_{BS} are consistent with a constant value (Figure 3). Quantitatively, we find that

$$\Pi_{\text{BS}} = 1350 \cdot \left[\frac{\langle F_{\text{max}} \rangle}{14.2} \right]^{0.03 \pm 0.03}, \quad (8)$$

where the units are the same as in equation (7). During the back slopes, the differences between average $E^{(i)}$ and average $\tau_{\text{BS}}^{(i)}$ (Table 1) for different intensity groups $i = 1\text{--}6$ effectively compensate each other when they form such a product as Π_{BS} . Since the

two are physically different parameters of emitters, we conclude that their brightness dependencies have predominantly a geometrical origin, i.e., they are due to the geometrical transformations of time and energy in the expanding Universe.

Comparing the values of $\langle E_p \rangle$ and τ_{BS} for the 100 brightest bursts (group 1, with peak fluxes $> 3.8 \text{ } \gamma/\text{cm}^2\text{-s}$) with those for the 100 dimmest bursts (group 6, with peak fluxes $< 0.43 \text{ } \gamma/\text{cm}^2\text{-s}$), we find that the factor of cosmological transformation between emitters of these groups, both for time dilation and red shift, is about 3 (Table 1). For this factor the value of z_{dim} for emitters of dimmest bursts is about 2, provided the brightest bursts correspond to $z_{\text{br}} \ll 1$. Recently it has been suggested that $z_{\text{br}} > 1$, based on measurements of the spectra of bright burst (Dezalay et al. 1998), and as the consequence of the idea that GRBs sources should follow the history of star formation (Wijers 1998, Che et al. 1998). In this case the group of dimmest bursts would have an average red-shift factor $z_{\text{dim}} \sim (3 \cdot (1 + z_{\text{br}}) - 1)$ as large as ~ 5 . While the outbursting sources are effectively standard candles along the back slopes, they are not standard along the rise fronts of bursts. If the bursts have a cosmological origin, observations of dim and bright bursts correspond in local time to the younger and older Universe, respectively. The variation of Π_{RF} with intensity (Table 2) is associated with a difference of average duration τ_{RF} in the co-moving frames of reference, because the values of $E_{p,\text{max}}$ are the same for both Π_{BS} and Π_{RF} . In the co-moving frames the bursts in the recent Universe have an average rise time ~ 3 times shorter than the average rise time of bursts in the early Universe.

The difference in rise time between emitters of bright and dim bursts could be the result of differences in the interaction of the outbursting source with the surrounding medium. Emitters of dimmer bursts could have interacted in a medium with higher density, or with a harder background emission, or with a stronger average magnetic field, or with a difference of some other global parameter of the Universe. On the other hand, during

the back slope there is no difference between bursts from close and distant cosmological distances. The tails of bursts are thought to represent a self-determined internal process, which has some internal time scale associated either with some inertia, or with a time constant of some decay, or with some other process, and which does not depend on the external condition of the surrounding medium. A future cosmological model has to take into account these differences between processes of emission during the rising phases of bursts and their decays, which is apparently intrinsic to the co-moving reference frames.

Fig. 1.— The $E_{\text{p,max}}$ distributions for different brightness groups. The distribution for the reference group is showed by thick line. The distribution for the dimmest group (6) is showed by thin line.

Fig. 2.— The dependence of $ACIP_{\text{RF}}$ on $\langle F_{\text{max}} \rangle$. The best linear fit (solid curve) has a non-zero slope.

Fig. 3.— The dependence of $ACIP_{\text{BS}}$ on $\langle F_{\text{max}} \rangle$. The best linear fit (solid curve) has a slope consistent with zero.

REFERENCES

- Anfimov, D. S., et al., 1998, in Gamma-Ray Bursts: 4th Huntsville Symposium, ed. C. A. Meegan, R. D. Preece & T. M. Koshut (New York: AIP), 289
- Brainerd, J. J. 1997, ApJ, 487, 96
- Che H. Yang Y. Nemiroff R.J. 1998, astro-ph/9810220
- Dezalay J-P., et al., 1998, in Gamma-Ray Bursts: 4th Huntsville Symposium, ed. C. A. Meegan, R. D. Preece & T. M. Koshut (New York: AIP), 15
- Kommers, J. M., Lewin, W. H. G., Kouveliotou, C., van Paradijs, J., Pendleton, G. N., Meegan, C. A., & Fishman, G. J., 1997, ApJ, 491, 704.
- Kulkarni, S. R., et al. 1998, Nature, 393, 35
- Litvak, M. L., et al. 1998, in Gamma-Ray Bursts: 4th Huntsville Symposium, ed. C. A. Meegan, R. D. Preece & T. M. Koshut (New York: AIP), 256
- Mallozzi, R. S., et al. 1995, ApJ, 454, 597
- Meegan, C. A., et al. 1992, Nature, 355, 143
- Meegan, C. A., et al. 1996, ApJS, 106, 65
- Metzger, M. R., et al. 1997, Nature, 387, 878
- Mitrofanov, I. G. 1996, in Gamma-Ray Bursts: 3rd Huntsville Symposium, ed. C. Kouveliotou, M. S. Briggs & G. J. Fishman (New York: AIP), 3
- Mitrofanov, I. G., et al. 1996, ApJ, 459, 570
- Mitrofanov, I. G., et al. 1997b, ApJ, 490, 509

Mitrofanov, I. G., et al. 1997, ApJ, 474, 272

Mitrofanov, I. G., et al. 1998, ApJ, in press

Mitrofanov I. G., et al., 1998a, in Gamma-Ray Bursts: 4th Huntsville Symposium,
ed. C. A. Meegan, R. D. Preece & T. M. Koshut (New York: AIP), 20

Norris, J. P., et al. 1994, ApJ, 424, 540

Norris, J. P., et al. 1998, in Gamma-Ray Bursts: 4th Huntsville Symposium,
ed. C. A. Meegan, R. D. Preece & T. M. Koshut (New York: AIP), 166

Paczynski, B. 1986, ApJ, 308, L5

Wijers, R.A.M.J. and Paczynski B. 1994, ApJ, 437, L107

Wijers, R.A.M.J. et al. 1998, MNRAS, in press

Table 1.

Intensity group	Peak flux ($\gamma/\text{cm}^2\text{s}$)	Average peak flux ($\gamma/\text{cm}^2\text{s}$)	$\langle E_p \rangle$ (keV)	Y	τ_{RF} (s)	τ_{BS} (s)
1	> 3.8	14.2 ± 1.4	293 ± 23	1.0	4.1 ± 0.5	4.6 ± 0.6
2	1.6-3.8	2.4 ± 0.3	235 ± 21	1.22 ± 0.12	4.7 ± 0.6	7.3 ± 0.9
3	0.95-1.6	1.2 ± 0.1	160 ± 12	1.82 ± 0.35	4.9 ± 0.6	8.8 ± 1.1
4	0.62-0.95	0.77 ± 0.08	134 ± 14	2.07 ± 0.20	4.6 ± 0.6	8.0 ± 1.0
5	0.43-0.62	0.51 ± 0.05	116 ± 12	2.58 ± 0.23	5.3 ± 0.7	10.2 ± 1.3
6	< 0.43	0.33 ± 0.03	97 ± 10	2.92 ± 0.10	4.5 ± 0.6	12.4 ± 1.6

Table 2.

Intensity group	Π_{RF} (keV-s)	Π_{BS} (keV-s)
1	1196 ± 174	1350 ± 205
2	1107 ± 172	1716 ± 271
3	784 ± 113	1402 ± 211
4	616 ± 103	1072 ± 179
5	613 ± 103	1183 ± 194
6	438 ± 74	1203 ± 199





